

- Baroni, C., and Hall, B. L. (2004). A new relative sea-level curve for Terra Nova Bay, Victoria Land, Antarctica. *Journal of Quaternary Science* 19, 377–396.
- Bassett, S. E., Milne, G. A., Mitrovica, J. X., and Clark, P. U. (2005). Ice sheet and solid earth influences on far-field sea-level histories. *Science* 309, 925–928.
- Bentley, M. J., Hodgson, D. A., Smith, J. A., and Cox, N. J. (2005). Relative sea level curves for the South Shetland Islands and Marguerite Bay, Antarctic Peninsula. *Quaternary Science Reviews* 24, 1203–1216.
- Blake, W., Jr. (1970). Studies of glacial history in Arctic Canada. *Canadian Journal of Earth Sciences* 7, 634–664.
- Blake, W., Jr. (1975). Radiocarbon age determination and post-glacial emergence at Cape Storm, southern Ellesmere Island. *Geografiska Annaler* 57A, 1–71.
- Clark, J. A., Farrell, W. E., and Peltier, W. R. (1978). Global changes in postglacial sea level: a numerical calculation. *Quaternary Research* 9, 265–287.
- Clark, P. U., and Mix, A. C. (2002). Ice sheets and sea level of the Last Glacial Maximum. *Quaternary Science Reviews* 21, 1–7.
- Conway, H., Hall, B. L., Denton, G. H., Gades, A. M., and Waddington, E. D. (1999). Past and future grounding-line retreat of the West Antarctic Ice Sheet. *Science* 286, 280–283.
- Dyke, A. S. (1998). Holocene delevelling of Devon Island, Arctic Canada: implications for ice sheet geometry and crustal response. *Canadian Journal of Earth Sciences* 35, 885–904.
- Dyke, A. S. (1999). The last glacial maximum and the deglaciation of Devon Island: support for an Innuitian Ice Sheet. *Quaternary Science Reviews* 18, 393–420.
- England, J. (1976). Late Quaternary glaciation of the Queen Elizabeth Islands, Northwest Territories, Canada: alternative models. *Quaternary Research* 6, 185–202.
- England, J. (1983). Isostatic adjustments in a full glacial sea. *Canadian Journal of Earth Sciences* 20, 895–917.
- England, J. (1992). Postglacial emergence in the Canadian High Arctic: integrating glacioisostasy, eustasy, and late deglaciation. *Canadian Journal of Earth Sciences* 29, 984–999.
- England, J. (1998). Support for the Innuitian Ice Sheet in the Canadian High Arctic during the Last Glacial Maximum. *Journal of Quaternary Science* 13, 275–280.
- England, J. (1999). Coalescent Greenland and Innuitian ice during the Last Glacial Maximum: revising the Quaternary of the Canadian High Arctic. *Quaternary Science Reviews* 18, 421–556.
- England, J., Atkinson, N., Bednarski, J., Dyke, A. S., Hodgson, D. A., and Ó Cofaigh, C. (2006). The Innuitian Ice Sheet: configuration, dynamics and chronology. *Quaternary Science Reviews* 25, 689–703.
- Hall, B. L., and Denton, G. H. (1999). New relative sea-level curves for the southern Scott Coast, Antarctica: evidence for Holocene deglaciation of the western Ross Sea. *Journal of Quaternary Science* 14, 641–650.
- Hall, B. L., Baroni, C., and Denton, G. H. (2004). Holocene relative sea-level history of the southern Victoria Land Coast, Antarctica. *Global and Planetary Change* 42, 241–263.
- Hodgson, D. A., Taylor, R. B., and Fyles, J. G. (1994). Late Quaternary sea level changes on Brock and Prince Patrick islands, western Canadian Arctic Archipelago. *Géographie physique et Quaternaire* 48, 69–84.
- Long, A. J., Roberts, D. H., and Wright, M. (1999). Isolation basin stratigraphy and Holocene relative sea-level change on Arveprinsen Eiland, Disko Bugt, West Greenland. *Journal of Quaternary Science* 14, 323–345.
- Ó Cofaigh, C. (1999). Holocene emergence and shoreline delevelling, southern Eureka Sound, High Arctic Canada. *Géographie physique et Quaternaire* 53, 235–247.
- Ó Cofaigh, C., England, J., and Zreda, M. (2000). Late Wisconsinan glaciation of southern Eureka Sound: evidence for extensive Innuitian ice in the Canadian High Arctic during the Last Glacial Maximum. *Quaternary Science Reviews* 19, 1319–1341.
- Quinlan, G., and Beaumont, C. (1981). A comparison of observed and theoretical postglacial relative sea level in Atlantic Canada. *Canadian Journal of Earth Sciences* 18, 1146–1163.
- Walcott, R. I. (1972). Late Quaternary vertical movements in eastern North America: quantitative evidence of glacio-isostatic rebound. *Reviews of Geophysics and Space Physics* 10, 849–884.
- Zwartz, D., Bird, M., Stone, J., and Lambeck, K. (1998). Holocene sea level and ice-sheet history in the Vestfold Hills, East Antarctica. *Earth and Planetary Science Letters* 155, 131–145.


## Mid-Latitudes

B P Horton, University of Pennsylvania, PA, USA

© 2007 Elsevier B.V. All rights reserved.

## Introduction

Since the Last Glacial Maximum (LGM), approximately 50 million cubic kilometers of ice melted from the land-based ice sheets (*see* Eustatic Sea-Level Changes Since the Last Glaciation), raising global relative sea level (RSL) in regions distant from the major glaciation centers (far-field sites; *see* Tropics) ca. 120 m (Fig. 1). In contrast, RSLs have dropped many hundreds of meters in regions once covered by the major ice sheets (near and intermediate field sites; *see* High Latitudes) as a consequence of the isostatic rebound of the solid Earth. Such rapid changes in RSL are part of a complex pattern of interactions among eustatic, isostatic (glacio and hydro), and local factors, all of which have different response timescales. The eustatic contribution to RSL change during deglaciation averaged 10 mm per year; however, peak rates potentially exceeded 50 mm per year during ‘meltwater pulses’ at 19 and 14.5 cal kyr BP (*see* Eustatic Sea-Level Changes Since the Last Glaciation). Empirical and glacial isostatic modeling studies suggested a significant reduction in the eustatic contributions to RSL change at ca. 7 cal kyr BP and the Earth entered into a period of RSL stability during which ocean volume, on average, changed only by a few meters. Clarke *et al.* (1978) identified six types of sea-level curve (I–VI), reflecting a range of RSL histories recorded in coasts which have emerged, submerged, or are in transitional areas, and record a combination of both uplift and subsidence (*see* High Latitudes, Tropics). Although these curves provide the general impression



**Figure 1** The relative sea-level curve for the last glacial cycle for Huon Peninsula supplemented with observations from Bonaparte Gulf, Australia. Error bars define the upper and lower limits. Modified from Lambeck K, Esat TM, and Potter E-K (2002) Links between climate and sea levels for the past three million years. *Nature* 419: 199–206.


of the rate and direction of RSL change, they do not reflect the true uncertainty associated with estimates of the altitude and age of former sea levels (Fig. 2). This article examines the processes and patterns of late Quaternary sea-level changes along the passive coastal margins of the mid-latitudes, with reference to selected studies. The article includes all types of sea-level curve except Zone I (Clarke *et al.*, 1978).

## Late Quaternary Sea Levels

### Atlantic Seaboard of North America

Atlantic Canada and the Gulf of Maine have had a complex, regionally and temporally varying sea level

since the LGM and represent Transitional Zones I–II and Zone II (Clark *et al.*, 1978). Shaw *et al.* (2002) presented the paleogeography of Atlantic Canada from 13  $^{14}\text{C}$  kyr BP to the present based upon the collection and compilation of sea-level index points from numerous studies. RSL curves from Newfoundland showed either continuous falling RSL with a marine limit at 120 m (e.g., Pinwar; Fig. 3A) or RSL dropping below the modern level before rising once more (e.g., Port au Port). Submerged Holocene deltas recorded the spatially and temporally varying postglacial RSL lowstand along the coast of Newfoundland. For example, RSL at La Poile Bay fell to a  $-30$  m lowstand ca. 10  $^{14}\text{C}$  kyr BP. The RSL curve for Nova Scotia falls to a lowstand of  $-65$  m at 11.3–11.7  $^{14}\text{C}$  kyr BP, before rising at a decreasing rate through the Holocene.




**Figure 2** Sea-level zones and typical relative sea-level curves deduced for each zone by Clark *et al.* (1978) under the assumption that no eustatic change has occurred since 5 kyr BP.

Since the LGM, isostatic rebound has dominated the RSL history of the northwestern Gulf of Maine. The retreat of the Laurentide ice sheet exposed large areas of isostatically depressed land, which were rapidly submerged by the sea (e.g., Barnhardt *et al.* 1995). The maximum inland extent of the marine invasion was 70–129 m above present sea level at 14  $^{14}\text{C}$  kyr BP (Fig. 3B). After ice retreat, the land rebounded isostatically, resulting in an RSL fall to a lowstand of  $-60$  to  $-65$  m on the inner shelf at 10.5  $^{14}\text{C}$  kyr BP. Lowstand shorelines and deltas recorded this phase. Subsequent slowing of the rate of rebound and overtaking by the rise of eustatic contributions resulted in local RSL rise at a generally decreasing rate to present. Sea-level rise over the past 5 cal kyr BP is known in considerable detail from high-marsh peats (*see* Low Energy Coasts Sedimentary Indicators, Microfossil Reconstructions). For example, Gehrels *et al.* (2004) showed a northeast to southwest gradient of crustal motion, reflecting the former distribution of Laurentide ice, from late Holocene RSL data from the Canadian provinces of New Brunswick and Nova Scotia, and the Gulf of Maine. The highest rates of late Holocene RSL rise occurred in Nova Scotia (ca. 2.5 mm per year). The RSL history for the southern

New Brunswick showed a late Holocene RSL rise of ca. 1 mm per year.

The Atlantic coast of the United States, from New Hampshire to Florida, has a wide continental shelf and great interplay between postglacial isostatic recovery, forebulge collapse, and hydroisostatic loading. Clark *et al.* (1978) suggested that the sea-level observations should be continually rising through the Holocene (Zone II), although records from Florida are supposed to show former sea levels above present (Zone III). Along the New Hampshire coast, saltmarsh deposits and drowned forests revealed that RSL has risen since 7  $^{14}\text{C}$  kyr BP, though at a rate decreasing gradually since ca. 4  $^{14}\text{C}$  kyr BP (Pirazzoli, 1991).

At ca. 13.5 cal kyr BP, RSL for Massachusetts was  $-65$  m below present (Uchupi *et al.*, 2006). RSL rose rapidly to 4.5 cal kyr BP, with a more gradual incline to near its present level ca. 1 cal kyr BP. In the Boston area of Massachusetts, there was a very limited rate of rise (0.6 m) for the last 3  $^{14}\text{C}$  kyr BP, with a more rapid rise (3 mm per yr) during the mid-Holocene (e.g., Pirazzoli, 1991). Donnelly (*in press*) studied basal high-marsh sediments from Romney Marsh, Revere, Massachusetts. He indicated a rise in RSL of ca. 2.6 m in the last 3.3 cal kyr. The data suggested a possible decrease in the rate of rise from 0.8 mm per year



**Figure 3** Selected relative sea-level observations for the mid-latitude: (A) Newfoundland (Shaw *et al.*, 2002); (B) Maine (Barnhardt *et al.*, 1995); (C) Delaware (Nikitina *et al.*, 2000), which includes wide error envelope; (D) Estonia (Raukas, 2000); (E) northwest Scotland (Shennan and Horton, 2002), which includes index points and limiting points (data points that are above RSL); (F) Belgium (Denys and Baetman, 1995) including index points and an RSL error band; (G) Portugal (Boski *et al.*, 2002); (H) south Australia (Harvey 2003); (I) South Africa (Compton, 2001); and (J) Beagle Channel (Milne *et al.*, 2005). Clarke's *et al.*'s (1978) six types of sea level curve (I–VI) are shown.

between 3.3 and 1 cal kyr BP to 0.5 mm per yr between 1 cal kyr BP and the last 150 to 500 yr. In southeastern Massachusetts, radiocarbon dates from shells and freshwater peat suggested that RSL rose from ca.  $-70$  m at 12  $^{14}\text{C}$  kyr BP to ca.  $-35$  m at 10  $^{14}\text{C}$  kyr BP at a rapid rate of 17 mm per year. Between 10  $^{14}\text{C}$  and 6  $^{14}\text{C}$  kyr BP, the rate of RSL rise dropped to ca. 3 mm per year and remained at that rate until ca. 2  $^{14}\text{C}$  kyr BP. During the last 2  $^{14}\text{C}$  kyr BP, the general rate of RSL rise has been ca. 1 mm per year (Pirazzoli, 1991).

RSL changes in Connecticut have been inferred from New Haven Harbor and Hammock River marshes. The former site revealed an RSL rise of 10.5–12 m during the last 5.9  $^{14}\text{C}$  kyr BP, whereas the latter did not show any pauses or reverses in the rise of RSL for the last 7  $^{14}\text{C}$  kyr BP. Van de Plassche *et al.* (1998) created a new RSL curve from the Hammock River marsh for the last 1.5 cal kyr using AMS radiocarbon dates from salt marsh deposits. The resulting record, expressed as deviations from

a long-term (isostatic) trend of 1 mm per year, suggested that sea level has oscillated centimeters to decimeters on timescales of centuries, being 0.25 m higher during the Medieval Warm Period than the Little Ice Age and rising at 1 mm per year over the last 300 cal yr. Donnelly *et al.* (2004) constructed a high-resolution RSL record for the past 700 yr by dating basal salt-marsh peat samples above a glacial erratic from an eastern Connecticut salt marsh. The data revealed an RSL rise of 1 mm per year from AD 1300 to 1850 with a nearly threefold increase in the regional rate of sea-level rise to modern levels that likely occurred in the later half of the nineteenth century.

For the New York region, RSL since the LGM was amplified by the proposed catastrophic drainage of late Wisconsin glacial lakes 12–14 cal kyr BP (e.g., Theiler *et al.*, in press). The shoreline was located at ca. 60 m isobath and the large glacial lakes north of the New York Bight were thought to have breached the moraine front at the Verrazano Narrows

and other locations in New Jersey and New York. Gornitz and Seeber (1990) suggested a 2 mm per year RSL rise during the last 7  $^{14}\text{C}$  kyr BP. Sea-level rise in the last 150 yr appears to have exceeded historic rates. Recent sea-level studies of the New Jersey coast suggested a rise of 2 mm per year from 6 cal kyr BP (e.g., Miller *et al.*, 2005). The Delaware RSL curve suggested rising RSL rates of 3 mm per year before 5 cal kyr BP, 2.1 mm per year from 5 to 2 cal kyr BP and 0.9 mm per year from 1.3 cal kyr BP to present (Nikitina *et al.*, 2000; Fig. 3C). Chesapeake Bay, the largest estuary in the US, is a geologic product of rising sea level. By 10 cal kyr BP, the main channel of the ancient Susquehanna River valley was flooded and became a narrow estuary (e.g., Larsen, 1998). Between 6 and 7 cal kyr BP, RSL was ca. 9 m below present and Chesapeake Bay took on its characteristic 'drowned river valley' shoreline pattern. The rate of RSL rise from 6 to 1 cal kyr BP was 1.4 mm per year. The rate further decreased to 0.6 mm per year in the last 1 kyr (Kearney, 1996).

There is a shortage of RSL data along the Atlantic Coast from the Carolinas to Florida at the boundary between Zones II and III (Clark *et al.*, 1978). In South Carolina, an oscillating RSL history during the Holocene has been proposed (e.g., Scott *et al.*, 1995). RSL rose from  $-3$  m at 4.6–5.2 cal kyr BP to  $+1$  m by ca. 4.3 cal kyr BP. Sea level subsequently fell to  $-3$  m by ca. 3.6 cal kyr BP. Since 3.6 cal kyr BP, sea level has risen steadily at a rate of 1 mm per year. A few notable papers from the Florida Keys showed continual rise of RSL during the Holocene with no indication of an emergence in Florida during the Holocene (e.g., Toscano and MacIntyre, 2003). Using U–Th dating of pristine *Acropora palmata* and head corals cored, Toscano and MacIntyre (2003) suggested RSL rose from  $-13.5$  to  $-7$  m between 8.9 and 5 ka. Toscano and MacIntyre (2003) identified a catastrophic RSL rise event of  $>45$  mm per year, between 7.6 and 7.2 ka, which they attributed to west Antarctic ice sheet instability and changes in marine ice extent between 8 and 7 ka. Sea-level reconstructions based on soilstones and mangrove peat showed a rise of 1.2 mm per year from  $-7$  m at 7  $^{14}\text{C}$  kyr BP to  $-0.8$  m at 2  $^{14}\text{C}$  kyr BP, followed by a rise of 0.3 mm per year from 2 cal kyr BP to the present.

## Europe

RSL observations and geophysical isostatic models clearly define the area of postglacial rebound in Fennoscandia (*see* Isostasy). This is surrounded by a subsiding zone that has the greatest postglacial subsidence (the so-called glacial forebulge or peripheral

bulge), which is situated in the North Sea between Norway and Great Britain, and extends through the northwestern Netherlands and northern Germany. Clark *et al.* (1978) suggest European RSL observations should represent Transitional Zone I–II, Zone II and Zone III.

Late Quaternary RSL change has led to the evolution of coastlines along the southern Baltic Sea. The Baltic Ice Lake (the first stage in the history of the Baltic Sea) formed 12  $^{14}\text{C}$  kyr BP after the readvance of the ice margin in north Estonia (Raukas, 2000). The retreat of the continental ice cover in central Sweden at 10.6  $^{14}\text{C}$  kyr BP caused the lowering of the Baltic Ice Lake level by 25–30 m. Sea level rapidly rose to 42–45 m above present at 9.2–9  $^{14}\text{C}$  kyr BP on the Island of Hiiumaa in Estonia (Raukas, 2000; Fig. 3D). Sea level subsequently fell 30–35 m between 9 and 8  $^{14}\text{C}$  kyr BP. Eustatic contributions to sea level exceeded the rate of glacioisostatic land uplift in the eastern part of the Gulf of Finland in the period 8.4–6.8 cal kyr BP. In southeastern Sweden, several minor fluctuations of RSL were identified between 8.5 and 5 cal kyr BP (Berglund *et al.*, 2005). They correlated a distinct regression phase around 8.1 cal kyr BP with the Greenland ice-core cold event dated to 8.2 ice-core kyr BP. There was a rapid RSL fall from the LGM in southern Sweden until approximately 9.2  $^{14}\text{C}$  kyr BP followed by RSL rise with a series of observations until 7.2  $^{14}\text{C}$  kyr BP and then a fall in RSL to present. The uplift rate has been estimated at 2.5 mm per year during the last 7 kyr. In Baltic Germany, RSL rose rapidly (20 mm per year) from 8 to 5.7 cal kyr BP, after which RSL gradually increased to present (e.g., Meyer and Harff, 2005). Danish coastlines show evidence of emergence in the north and submergence to the south because it is located at the margin of the former Scandinavian ice sheet. For example, Clemmensen *et al.* (2001) produced a Holocene RSL curve for Skagen at the northern tip of Denmark. Sea level fell ca. 11 m between 5.6 and 1.4 cal kyr BP, but was characterized by periods of stability (ca. 5.7–5.5 cal kyr BP, 5.2–4.7 cal kyr BP, 4.1–2.6 cal kyr BP, and 1.7–1.5 cal kyr BP) separated by periods of rapid fall.

Shennan and Horton (2002) provided a synthesis of several local and regional RSL analyses and proposed simplified late Quaternary summary RSL curves for regions of the United Kingdom. Isolation basins, raised tidal marshes, coastal wetlands and dune systems from northwest Scotland have produced a 16 cal kyr BP record of RSL change from the time of local deglaciation following the LGM to the present (Fig. 3E). Sea-level records from Scotland reflected the spatially variable effect of glacioisostatic adjustment, which produced a mid-Holocene

highstand that diminishes from maximum altitudes at the sites beneath thickest ice at LGM to the peripheral sites under thin ice cover, with the Shetland Isles showing no Holocene RSL above present. The transition at the north appeared to be between Orkney and Wick, which showed a small highstand. On both the east and west coasts of the UK, there was a clear north-to-south trend. For example, there was a regional difference between Northumberland and Norfolk, east coast of England, of 20 m at 8 cal kyr BP and by 4 cal kyr BP. RSL in Northumberland was above present, whereas in areas to the south RSL has been below present throughout the Holocene. Estimates for late Holocene RSL change ranged from 0.7 mm per year in the Norfolk to  $-0.7$  mm per year in Northumberland (Shennan and Horton, 2002).

During the LGM the southern North Sea was 110–130 m below present sea level, so most parts of the North Sea basin were dry land (e.g., Streif, 2004). Marine mollusks found at a depth of 72 m below present-day sea level yielded a radiocarbon age of 10.3  $^{14}\text{C}$  kyr BP. A phase of the transgression lasted from 10.3 to 7.1  $^{14}\text{C}$  kyr BP when RSL rose from 72 to 25 m below present (Streif, 2004). The average rate of RSL rise was ca. 15 mm per year, but reached ca. 21 mm per year between 8.6 and 7.1  $^{14}\text{C}$  kyr BP. The final phase of the transgression began at ca. 7.5  $^{14}\text{C}$  kyr BP and has continued until today. However, Behre (2004) suggested that RSL in the late Holocene was not continuous with a short phase of lowering in the thirteenth and fourteenth centuries.

Denys and Baeteman's (1995) Holocene sea-level data for the Belgian coastal plain suggested an initial RSL rise of ca. 7 mm per year before ca. 7.5 cal kyr BP. This resulted in a very rapid shift of the facies belts across the continental shelf toward a position close to the present-day boundary of the coastal plain. At ca. 7.5–7 cal kyr BP, the RSL curve showed a distinct rate-of-rise decrease to an average of 2.5 mm per year, and consequently, the rapid landward shift of the various sedimentary environments ceased. The rate of RSL rise continued to decrease, and after ca. 5.5–5 cal kyr BP, it fell to an average of 0.7 mm per year. Van de Plassche (1982), who used predominantly basal peat data, concluded that during the past 6 cal kyr BP the northern and western Netherlands showed continuous RSL rise with an increase of over 1 m during the last 2.5 cal kyr BP that included a few small fluctuations.

Sea-level studies from Atlantic France showed a rapid rise in the early Holocene with no evidence of RSL being above present (Louwyte and Declercq, 1998). In the Gironde Estuary, Mellalieu *et al.*, (2000) identified a widespread deposit of fresh and brackish water peat that began forming across much

of the area from ca. 6.4 cal kyr BP. This peat reached thicknesses of 1–3 m and was inundated by marine conditions once more at ca. 2.8 cal kyr BP.

Portuguese RSL data are characteristic of Zone III (Clark *et al.*, 1978). RSL was ca. 100 m below present at 16 cal kyr BP. Since 13 cal kyr BP, a very rapid RSL occurred with RSL reaching  $-40$  m between 12 and 11 cal kyr BP followed by an equally rapid descent to  $-60$  m in response to the Younger Dryas (Dias *et al.*, 2000). Sea level rose rapidly again at ca. 10 cal kyr BP. Boski *et al.* (2002) indicated RSL rising rapidly at a rate of 8.5 mm per year to approximately 6.5 cal kyr BP (Fig. 3G). Since then, lagoonal sediments in the vicinity of the estuary have been enclosed behind sand spits and predominantly sandy sedimentation was initiated within the estuary. After a second phase of slower rise at the rate of 3 mm per year, which lasted until ca. 5 cal kyr BP, the sea approached the present level. There was no indication of an emergence during the Holocene.

### Southern Hemisphere

The available mid-latitude RSL observations from the Southern Hemisphere fall within Zone V, and therefore should illustrate a mid-Holocene highstand (Clark *et al.*, 1978). Australia is relatively seismically stable with little evidence of tectonic deformation since the Mesozoic. It is situated at the center of the Australian–Indian plate and has passive continental margins. This makes Australia an important far-field location for RSL study, ranging from the first widely proposed eustatic sea-level curve (*see* Eustatic Sea-Level Changes Since the Last Glaciation), to some of the earliest debates on hydroisostasy (*see* Isostasy).

In New South Wales, RSL rose rapidly from ca.  $-120$  m at the LGM to  $+1$  m by 6.5  $^{14}\text{C}$  kyr BP (Sloss *et al.*, 2005). In South Australia and Tasmania, RSL had risen sufficiently by 17.5  $^{14}\text{C}$  kyr BP to enter the Bass Basin from the west and form an estuarine environment (Lambeck and Chappell, 2001). At 14  $^{14}\text{C}$  kyr BP, the RSL rise reached the barrier in the east and Tasmania became isolated from the Australian mainland. Detailed geological investigations in South Australia reveal rapid rates of sea-level rise of 9 mm per year between 10 and 8  $^{14}\text{C}$  kyr BP (Harvey (2003), Fig. 3H), and a more rapid rate of 24 mm per year between 8 and 6.7  $^{14}\text{C}$  kyr BP. The peak of the postglacial marine transgression was  $+4.5$  m at ca. 7  $^{14}\text{C}$  kyr BP. Sites on the south coast of Western Australia displayed an RSL of  $+1$  m by ca. 7.5 cal kyr BP. There appears to have been a substantial stepped fall of RSL on the western Australian coast somewhere between 3.8 cal kyr BP and after 3.6 cal kyr BP. This is not what would be

expected from a dominant hydroisostatic signal, and is more suggestive of eustatic contributions (Lessa and Masselink, 2006).

The most complete and currently accepted Holocene RSL curve for New Zealand is largely based on data from tectonically 'quiet' areas near Dunedin and Auckland (e.g., Hayward *et al.*, 2002). Sea level rose from ca. 25 m below present at ca. 9 cal kyr BP to reach its present height at ca. 7 cal kyr BP. Since then, RSL has been no more than 1 m above and no more than 0.5 m below the present height.

Being remote from Quaternary ice sheets, the South African coasts have experienced only minor isostatic movements. Ramsay (1995) produced a 9-kyr record showing early Holocene RSL rise to a mid-Holocene highstand of +3.5 m at 4.7 <sup>14</sup>C kyr BP with RSL subsequently falling below present levels, but also showed a secondary highstand at 1.6 <sup>14</sup>C kyr BP (+1.5 m) before current RSL is attained at 0.9 <sup>14</sup>C kyr BP (Ramsay, 1995). A second investigation in South Africa has concentrated on a salt-marsh lagoon on the southwest coast (Compton, 2001). This lagoon reflected more Atlantic rather than Indian Ocean processes. The RSL record,

derived from calibrated radiocarbon-dated saltmarsh peats, agreed with Ramsay's two mid-late Holocene highstands theory (Fig. 3I).

Milne *et al.* (2005) compiled Holocene RSL data from the Atlantic coast of South America to understand the cause of the observed spatial trend and estimate a eustatic signal for the Holocene. Milne *et al.* (2005) concluded that the quality of the RSL data from mid-Atlantic locations was poor, with large age and altitude uncertainties and added complications provided by the possible effects of tectonic and isostatic factors associated, respectively, with Andean uplift and the loading effects of the Patagonian and Antarctic ice complexes. Within the Beagle Channel, Milne *et al.* (2005) demonstrated a pronounced highstand of +6 m at 6 cal kyr BP, followed by a fall to present by 3 cal kyr BP (Fig. 3J).

## Conclusion

High-quality RSL data from the mid-latitudes reveal spatial and temporal variations among eustatic, isostatic (glacio and hydro), and local factors since the LGM. Such data are used for many applications,

**Table 1** Summary of relative sea-level change from selection mid-latitude regions

Region	Relative sea-level description	References
Newfoundland	From 13 <sup>14</sup> C kyr BP to present RSL curves show either continuous falling RSL with a marine limit at 120 m or RSL dropping below the modern level before rising once more	Shaw <i>et al.</i> (2002)
Marine	RSL fall from 70–129 m above present sea level at 14 <sup>14</sup> C kyr BP to a lowstand of –60 to –65 m at 10.5 <sup>14</sup> C kyr BP. RSL rise at a generally decreasing rate to present	Barnhardt <i>et al.</i> (1995)
Delaware	Continuous rising RSL from 12 cal kyr BP with rates of 3 mm per yr before 5 cal kyr BP, 2.1 mm per yr from 5 to 2 cal kyr BP and 0.9 mm per year from 1.3 cal kyr BP to present. There is no indication of an emergence during the Holocene	Nikitina <i>et al.</i> (2000)
Estonia	RSL fall from 12 <sup>14</sup> C kyr BP to present with a rapid rise to 42–45 m above present at 9.2–9 <sup>14</sup> C kyr BP. Sea level subsequently fell 30–35 m between 9 and 8 yr <sup>14</sup> C kyr BP	Raukas (2000)
Northwest Scotland	RSL fall from a marine limit between 35 and 40 m at ca. 16 kyr cal BP through to an early Holocene minimum. Sea-level index points define the culmination of the mid-Holocene RSL highstand of 7 m above present at 7.6–7.4 cal kyr BP. The RSL highstand persisted for more than 1 kyr before the onset of any significant RSL fall	Shennan and Horton (2002)
Belgium	Continuously rising RSL from 9.3 cal kyr BP to present with rates of 7 mm per year before 7.5 cal kyr BP. At 7.5–7 cal kyr BP the RSL curve shows a distinct rate-of-rise decrease to an average of 2.5 mm per year. After 5.5–5 cal kyr BP, RSL rise fell to an average of 0.7 mm per year. There is no indication of an emergence during the Holocene	Denys and Baetman (1995)
Portugal	Continually rising RSL from 9.8 cal kyr BP with no indication of an emergence during the Holocene. Sea level rose at a rate of 8.5 mm per year to 6.5 cal kyr BP. Subsequently RSL rose at a slower rate of 3 mm per year, which lasted until 5 cal kyr BP	Boski <i>et al.</i> (2002)
South Australia	RSL rise from 10 <sup>14</sup> C kyr to a mid-Holocene highstand of +4.5 m at ca. 7 <sup>14</sup> C kyr, and a subsequent fall to present. RSL rose at 9 mm per yr between 10–8 <sup>14</sup> C kyr BP, with a more rapid rate of 24 mm per year between 8–6. 7 <sup>14</sup> C kyr BP	Harvey (2003)
South Africa	RSL rise from –30 m between 10 and 9 cal kyr BP to 0–3 m above present by 6.8 cal kyr BP. Sea level was near present between 4.9 and 2.5 ka followed by a drop in sea level of 1–2 m between 2.5 and 1.8 ka. A brief sea-level highstand of approximately. 0.5–1.0 m at 1.3 cal kyr BP followed by a lowstand of around –0.5 to –1.0 m at ca. 0.7 cal kyr BP	Compton (2001)
Beagle Channel	RSL data demonstrate a pronounced highstand of +6 m at ca. 6 cal kyr BP, followed by a fall to present sea level by ca. 3 cal kyr BP	Milne <i>et al.</i> (2005)

ranging from calibrating models of Earth rheology and ice-sheet reconstructions to the development of coastal lowlands and human occupation. However, there are still many open questions related to reconstructing former RSL (see Low Energy Coasts Sedimentary Indicators, Microfossil Reconstructions). **Figure 3** and **Table 1** show the variety of methods of interpreting RSL changes. These RSL analyses attempt to determine how sea level has varied by plotting index points on a scatter plot with RSL as the dependent variable. However, sea-level index points should be shown with an error box or envelope that demonstrates the full assessment of RSL and age errors. Errors that are often ignored in sea-level analyses included: (1) the uncertainty in the relationship between a given indicator and the local to regional paleoenvironment in which it is formed (known as the indicative meaning); (2) sediment compaction and tidal range variations; and (3) calibration of radiocarbon dates, and if appropriate the application of the marine reservoir effect.

See also: **Sea Level Studies: Low Energy Coasts Sedimentary Indicators; Microfossil Reconstructions; Eustatic Sea-Level Changes Since the Last Glaciation; Isostasy. Sea-Levels, Late Quaternary: High Latitudes; Tropics.**

## References

- Barnhardt, W. A., Gehrels, W. R., Belknap, D. F., and Kelley, J. T. (1995). Late quaternary relative sea-level change in the western Gulf of Maine. Evidence for a migrating glacial forebulge. *Geology* **23**, 317–320.
- Behre, K.-E. (2004). Coastal development, sea-level change and settlement history during the later Holocene in the clay district of Lower Saxony (Niedersachsen), northern Germany. *Quaternary International* **112**, 37–53.
- Berglund, B. E., Sandgren, P., Barnekow, L., et al. (2005). Early Holocene history of the Baltic Sea, as reflected in coastal sediments in Blekinge, southeastern Sweden. *Quaternary International* **130**, 111–139.
- Boski, T., Moura, D., Camacho, S., et al. (2002). The postglacial sea-level rise along SE Portuguese Coasts as recorded in two estuaries. *Sedimentary Geology* **150**, 103–122.
- Clark, J. A., Farrell, W. E., and Peltier, W. R. (1978). Global changes in postglacial sea level a numerical calculation. *Quaternary Research* **9**, 265–287.
- Clemmensen, L. B., Richardt, N., and Andersen, C. (2001). Holocene sea-level variation and spit development: Data from Skagen Odde, Denmark. *The Holocene* **11**, 323–331.
- Compton, J. S. (2001). Holocene sea-level fluctuations inferred from the evolution of depositional environments of the southern Langebaan lagoon salt marsh, South Africa. *The Holocene* **11**, 395–405.
- Denys, L., and Baeteman, C. (1995). Holocene evolution of relative sea level and local mean high water spring tides in Belgium - A first assessment. *Marine Geology* **124**, 1–19.
- Dias, J. M. A., Boski, T., Rodrigues, A., and Magalhães, F. (2000). Coast line evolution in Portugal since the last glacial maximum until present - A synthesis. *Marine Geology* **170**, 177–186.
- Donnelly, J. P., (in press). A revised late Holocene sea-level record for northern Massachusetts, USA: *Journal of Coastal Research*.
- Donnelly, J. P., Cleary, P., Newby, P., and Ettinger, R. (2004). Coupling instrumental and geological records of sea-level change: Evidence from southern New England of an increase in the rate of sea-level rise in the late 19th century. *Geophysical Research Letters* **31**, 1–4.
- Gehrels, W. R., Milne, G. A., Kirby, J. R., Patterson, R. T., and Belknap, D. F. (2004). Late Holocene sea-level changes and isostatic crustal movements in Atlantic Canada. *Quaternary International* **120**, 79–89.
- Gornitz, V., and Seeber, L. (1990). Vertical crustal movements along the East Coast, North America, from historic and late Holocene sea level data. *Tectonophysics* **178**, 127–150.
- Harvey, N. (2003). Past rates of sea level change. *Pages News* **11**, 19–20.
- Hayward, B. W., Grenfell, H. R., Sandiford, A., Shane, P. A. R., Morley, M. S., and Alloway, B. V. (2002). Foraminiferal and molluscan evidence for the Holocene marine history of two breached maar lakes, Auckland, New Zealand. *New Zealand Journal of Geology and Geophysics* **45**, 467–480.
- Kearney, M. S. (1996). Sea level change during the last thousand years in Chesapeake Bay. *Journal of Coastal Research* **12**, 977–983.
- Lambeck, K., and Chappell, J. (2001). Sea level change through the last glacial cycle. *Science* **292**, 679–686.
- Lambeck, K., Esat, T. M., and Potter, E.-K. (2002). Links between climate and sea levels for the past three million years. *Nature* **419**, 199–206.
- Larsen, C. E., (1998). The Chesapeake Bay: Geologic Product of Rising Sea Level. US Geological Survey Fact Sheet 102–98.
- Lessa, G., and Masselink, G. (2006). Evidence of a Mid-Holocene sea level highstand from the sedimentary record of a macrotidal barrier and paleoestuary system in northwestern Australia. *Journal of Coastal Research* **22**, 100–112.
- Louwe, S., and Declercq, E. (1998). Relative water level change in the intracoastal zone of Belgium and northern France over the last 2500 years. *Boreas* **27**, 162–177.
- Mellalieu, S. J., Massé, L., Coquillas, D., Alfonso, S., and Tastet, J.-P. (2000). Holocene development of the east bank of the Gironde Estuary: Geoarchaeological investigation of the Saint Ciers-sur-Gironde marsh. In *Coastal and Estuarine Environments: Sedimentology, Geomorphology and Geoarchaeology* (K. Pye and J. R. L. Allen, Eds.), pp. 317–341. Geological Society, London.
- Meyer, M., and Harff, J. (2005). Modelling palaeo coastline changes of the Baltic Sea. *Journal of Coastal Research* **21**, 598–609.
- Miller, K. G., Kominz, M. A., Browning, J. V., et al. (2005). The Phanerozoic record of global sea-level change. *Science* **310**, 1293–1298.
- Milne, G. A., Long, A. J., and Bassett, S. E. (2005). Modelling Holocene relative sea-level observations from the Caribbean and South America. *Quaternary Science Reviews* **24**, 1183–1202.
- Nikitina, D. J., Pizzuto, J. E., Schwimmer, R. A., and Ramsey, K. W. (2000). An updated Holocene sea-level curve for the Delaware coast. *Marine Geology* **171**, 7–20.
- Pirazzoli, P. A. (1991). *Oceanography Series, vol. 58: World Atlas of Holocene Sea-level Changes*, p. 300. Elsevier, Amsterdam.

- Ramsay, P. J. (1995). 9000 years of sea-level change along the southern African coastline. *Quaternary International* 31, 71–75.
- Raukas, A. (2000). Rapid changes of the Estonian coast during the late-glacial and Holocene. *Marine Geology* 170(1), 21–27.
- Redfield, A. C. (1967). Postglacial change in sea level in the western North Atlantic Ocean. *Science* 157, 687–692.
- Scott, D. B., Gayes, P. T., and Collins, E. S. (1995). Mid-Holocene precedent for a future rise in sea level along the Atlantic coast of North America. *Journal of Coastal Research* 11, 615–622.
- Shaw, J., Gareau, P., and Courtney, R. C. (2002). Palaeogeography of Atlantic Canada. *Quaternary Science Reviews* 21(16–17), 1861–1878.
- Shennan, I., and Horton, B. P. (2002). Relative sea-level changes and crustal movements of the UK. *Journal of Quaternary Science* 16, 511–526.
- Sloss, C. R., Jones, B. G., Murray-Wallace, C. V., and McClennen, C. E. (2005). Holocene sea level fluctuations and the sedimentary evolution of a barrier estuary: Lake Illawarra, New South Wales, Australia. *Journal of Coastal Research* 2005 21, 943–959.
- Streif, H. (2004). Sedimentary record of Pleistocene and Holocene marine inundations along the North Sea coast of Lower Saxony, Germany. *Quaternary International* 112, 3–28.
- Thieler E, Butman B, Schwab W, *et al.* (in press) A catastrophic meltwater flood event and the formation of the Hudson shelf valley. *Palaeogeography, Palaeoclimatology, Palaeoecology*.
- Toscano, M. A., and MacIntyre, I. G. (2003). Corrected western Atlantic sea-level curve for the last 11,000 years based on calibrated  $^{14}\text{C}$  dates from *Acropora palmata* framework and intertidal mangrove peat. *Coral Reefs* 22, 257–270.
- Uchupi, E., Giese, G., Driscoll, N., and Aubrey, D. G. (2005). Postglacial geomorphic evolution of a segment of Cape Cod Bay and adjacent Cape Cod, Massachusetts, USA. *Journal of Coastal Research* 21, 1085–1106.
- van de Plassche, O. (1982). Sea-level change and water movements in the Netherlands during the Holocene. PhD Dissertation, Free University, Amsterdam.
- van de Plassche, O., van der Borg, K., and de Jong, A. F. M. (1998). Sea level-climate correlation during the past 1400 year. *Geology* 26(4), 319–322.

## Tectonic Locations

A R Nelson, US Geological Survey, CO, USA

© 2007 Elsevier B.V. All rights reserved.

### Introduction

Records of relative sea-level (RSL) change archive the response of the Earth's surface to tectonic processes during the Quaternary, the most recent period of Earth history. Landforms or deposits marking former shorelines, termed strandlines, are widespread datums that can be measured more precisely than other types of landforms and deposits produced in response to tectonic processes. Although limited to coasts, sequences of successive strandlines are commonly better preserved over

longer periods of time than other tectonic landforms. One-third to one-half of the Earth's marine coasts lie along or near tectonically active plate boundaries where the potential for large earthquakes threatens many population centers (Lajoie, 1986). Much of our understanding of tectonic processes over hundreds to hundreds of thousands of years has come from study of displacements (vertical components of tectonic deformation caused by faulting and local or regional folding) obtained through mapping and dating sequences of strandlines along tectonically active coasts. This improved understanding of the rates and scales of tectonic processes leads to better forecasting and mitigation of large earthquakes and related hazards—such as tsunamis and landslides—in inland as well as coastal regions worldwide.

RSL changes are a composite of real sea-level changes, which are vertical movements of the ocean's surface, and apparent sea-level changes, which are the inverse of vertical land-level changes along coasts (Lajoie, 1986; van de Plassche, 1986; Fig. 1). Although sea level may rise and fall by meters over periods of hours during tidal cycles, storm surges, or tsunamis, most long-term evidence of real changes in mean sea level consists of erosional or depositional strandlines formed over periods of years to millennia. Real changes recorded by strandlines range from local changes of decimeters, for example, caused by changes in the configuration of the mouth of an estuary following a severe storm, to fluctuations of global sea level of 100–150 m caused by expansion and contraction of continental ice sheets over tens of thousands of years (Chappell *et al.*, 1996). Causes of apparent sea-level changes range from the slow ( $<0.01 \text{ mm yr}^{-1}$ ) regional epeirogenic uplift of coasts over hundreds of thousands of years (Bloom, 1998), to the meters of instantaneous, localized uplift of some coasts during the largest subduction-zone (a regional fault where one tectonic plate slides beneath another) earthquakes (Berryman, 1987; Ota and Yamaguchi, 2004). Quantifying the tectonic displacement (uplift or subsidence) of strandlines requires separating the apparent changes from the real changes in strandline records.

The areal extent and scale of RSL changes, and the precision with which former sea levels can be measured from strandlines, depend on time span and sea-level history. For this reason, different types of information about tectonic processes are derived from different types of strandlines of differing age. Small apparent sea-level changes of a few meters or less are much more easily identified